

Global Ocean Prediction Using HYCOM

E.J. Metzger¹, A.J. Wallcraft¹, E.P. Chassignet², H.E. Hurlburt¹,
O.M. Smedstad³ and J.A. Cummings⁴

¹ Naval Research Laboratory, Stennis Space Center, MS

² Florida State University, Tallahassee, FL

³ Planning Systems Inc., Slidell, LA

⁴ Naval Research Laboratory, Monterey, CA

DoD HPC Modernization Program
Users Group Conference
26-29 June 2006, Denver CO


Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE JUN 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Global Ocean Prediction Using HYCOM				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,Stennis Space Center,MS,39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 25	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

HYCOM – HYbrid Coordinate Ocean Model

- Developed from MICOM by HYCOM/NOPP Consortium
 - Naval Research Lab, U. Miami, FSU, Los Alamos → GISS
 - Creating a true community ocean model
- Hybrid (generalized) vertical coordinate
 - Isopycnal in open, stratified ocean
 - Terrain-following in shallow coastal regions
 - Z-level in mixed layer and other unstratified regions
 - Generalized – not limited to these types
 - Dynamically smooth transition between coordinate systems
 - Coordinate choice varies in space and time
 - Isopycnals intersecting sloping topography by allowing zero thickness layers
 - Accurate transition between the deep and shallow water
- Existing mixed layer options
 - KPP Mellor-Yamada 2.5 GISS
 - Kraus-Turner Price-Weller-Pinkel

South China Sea Warm Current Feeding the Taiwan Current

Mean Zonal Velocity vs. Depth (top 500m) Along 117°E



HYCOM makes an accurate transition between the deep and coastal waters and can realistically simulate both

HYCOM Long-term Goals for Operational Ocean Prediction

- 1/12° fully global ocean prediction system (~7 km mid-latitude) transitioned to NAVOCEANO in 2007
 - Include shallow water, minimum depth 5 m
 - Coupled sea-ice model (Los Alamos CICE)
 - Data assimilation (NCODA)
- Increase to 1/25° resolution globally (~3-4 km mid-latitude) by the end of the decade
 - Optimal resolution for basin-scale
 - Boundary conditions for coastal models


Global HYCOM Configuration

- Horizontal grid: 1/12° equatorial resolution
 - 4500 x 3298 grid points, ~6.5 km spacing on average, ~3.5 km at pole
- Mercator 79°S to 47°N, then Arctic dipole patch
- Vertical coordinate surfaces: 32 for σ_2^*
- GISS mixed layer model
- Thermodynamic (energy loan) sea-ice model
- Surface forcing: wind stress, wind speed, thermal forcing, precipitation, relaxation to climatological SSS
- Monthly river runoff (986 rivers)
- Initialize from January climatology (GDEM3) T and S, then SSS relaxation from PHC 3.0
 - No subsurface relaxation to climatology

1/12° Global HYCOM

Sea surface height and ice (gray)


SSH year 7.75 (Oct 02)



- Running at NAVOCEANO on IBM Power 4+ (kraken)
- 216K CPU hrs/model year on 784 processors
- 3.1 Tb/model year for (compressed) daily 3-D output

Long-term Mean Global Sea Level


1992-2002 Mean Dynamic Ocean Topography (0.5°)



The 1992-2002 mean ocean dynamic topography data has been obtained from Nikolai Maximenko (IPRC) and Peter Niiler (SIO)

Long-term Mean Global Sea Level



1/12° global HYCOM



5 year model mean using climatological ECMWF wind and thermal forcing
HYCOM mean shifted by 10 cm

1/12° Global HYCOM

Mean Gulf Stream and Kuroshio pathways




Mean over five model years

ERA15 climatological wind &
thermal forcing

Sea Surface Height (SSH) Variability

Satellite altimetry (top) vs. 1/12° global HYCOM (bottom)




1992 – 2005 SSH variability based on T/P, ERS-1, and ERS-2 altimeters

SSH variability from 1/12° global HYCOM σ_2^* with climatological wind and thermal forcing


Global HYCOM is reproducing the expected eddy structure

Velocity Cross-section Along Luzon Strait

Sb-ADCP data (top) vs. 1/12° global HYCOM (bottom) in the upper 300 m
Section along 21°N between 118.5°E and 124.0°E




Cross-section overlaid on
mean currents and speed




Sb-ADCP data from Liang et al. (2003, DSR Pt. II)
Mean from HYCOM with ERA15 wind and thermal forcing
No ocean data assimilation in HYCOM

Velocity Cross-section East of Taiwan

Sb-ADCP data (top) vs. 1/12° global HYCOM (bottom) in the upper 300 m
Sections at 22°N (left) and 25°N (right), Taiwan coast to 124°E




Cross-section overlaid on mean currents and speed



Sb-ADCP data from Liang et al. (2003, DSR Pt. II)
Mean from HYCOM with ERA15 wind and thermal forcing
No ocean data assimilation in HYCOM

Velocity Cross-section Along the Equatorial Pacific

TOGA TAO data (left) vs. 1/12° global HYCOM (right) in the upper 400 m
Section between 143°E and 95°W




TOGA TAO data from Johnson et al. (2002, Prog. Oceanogr.)

5 year mean from HYCOM using high-frequency ECMWF winds and thermal forcing

No ocean data assimilation in HYCOM

Velocity Cross-section Across Equator at 140°W

TOGA TAO data (left) vs. 1/12° global HYCOM (right) in the upper 400 m
Section between 8°S and 10°N





TOGA TAO data from Johnson et al. (2002, Prog. Oceanogr.)

5 year mean from HYCOM using high-frequency ECMWF winds and thermal forcing

No ocean data assimilation in HYCOM

Velocity Cross-section Across Yucatan Channel

Mooring data (left) vs. 1/12° global HYCOM (right) in top 2200 m




From Candela et al. (2002, GRL)


Coastal/Island Sea Level Comparison From 1/12° Global HYCOM

2003 statistics at 126 tide gauge stations

RMS Difference





Correlation



Global HYCOM is reproducing the deterministic
response to the wind-driven circulation

SST Response in 1/12° Global HYCOM to Hurricanes Katrina and Rita



HYCOM reproduces the deterministic SST response to the wind forcing.
Implies realistic upwelling and mixing of subsurface waters as well
as realistic atmospheric wind and heat flux forcing in HYCOM.


One-way File-based Coupling Between HYCOM and CICE (PIPS 3.0)

Sea Ice Concentration (%) - September 2003

CICE stand-alone
(no ocean)


CICE with
HYCOM forcing

SSMI
observations



Too much ice


Coupling between the ocean and ice models more properly accounts for the momentum, heat and salt fluxes at the air/sea interface




ESMF-based Coupling Between .72° Arctic HYCOM and CICE

Sea Ice Thickness (m) and Drift - September 2003


File-based coupling



ESMF-based one-way coupling




NRL Coupled Ocean Data Assimilation (NCODA)



**MVOI - simultaneous analysis 5 ocean variables
temperature, salinity, pressure, velocity (u,v)**


Data Assimilation Subregions

Overlaid on SSH valid at 5 January 2004




Data Assimilation in Global HYCOM

Gulf Stream SSH with SST-based frontal analysis overlaid



← Initial state

After five assimilation cycles ↓





Frontal analysis < 4 days old = white,
analysis \geq 4 days old = black

Sea Surface Height Increments (top) and Observation Locations (bottom)

1 January 2004

5 January 2004



Lines: altimeter tracks; black dots: *in situ* observations

Future Plans

- FY06:
 - Final non-assimilative spin-up experiment (with improved wind/thermal forcing and CICE)
 - Multiple data-assimilative experiments over the period May 2001 – June 2002 to tune and refine the assimilation technique
 - May 2001 to present data-assimilative hindcast
 - Start a near real-time runs that mimic the expected operational procedure

Future Plans

- FY07:
 - Continue May 2001-June 2002 experiments, some with “advanced” assimilation
 - Complete a 1993-present “ocean reanalysis” by running a data-assimilative hindcast from 1993 up through May 2001
 - Two non-assimilative simulations:
 - 1995-2007 NOGAPS forcing
 - 1979-2006 ECMWF forcing
 - Ten year 1/25° Atlantic demonstration